

# Handy's Harbinger

David W. Schwenke  
NASA Ames Research Center  
Moffett Field, CA

Aug. 13, 2014

# Problem:

- ▶ The accurate computation of ro-vibrational energy levels of polyatomic molecules using potential energy functions
  - ▶  $\text{H}_2\text{O}$ , HCCH, ...

# Coordinates are King!

- ▶ Normal modes
  - ▶  $KE^1$  complicated,  $BC^2$  nasty
- ▶ Polyspherical Orthogonal coordinates
  - ▶ KE and BC simple, PES complicated
- ▶ Bond-Length-Bond-Angle coordinates (Polyspherical Non-orthogonal coordinates)
  - ▶ Quartic PES and BC simple
  - ▶ KE complicated

---

<sup>1</sup>Kinetic Energy Operator

<sup>2</sup>Boundary Conditions

# Coordinates are King!

- ▶ Normal modes
  - ▶  $KE^1$  complicated,  $BC^2$  nasty
- ▶ Polyspherical Orthogonal coordinates
  - ▶ KE and BC simple, PES complicated
- ▶ Bond-Length-Bond-Angle coordinates (Polyspherical Non-orthogonal coordinates)
  - ▶ Quartic PES and BC simple
  - ▶ KE complicated
- ▶ What is optimum?

---

<sup>1</sup>Kinetic Energy Operator

<sup>2</sup>Boundary Conditions

# Coordinates are King!

- ▶ Normal modes
  - ▶ KE<sup>1</sup> complicated, BC<sup>2</sup> nasty
- ▶ Polyspherical Orthogonal coordinates
  - ▶ KE and BC simple, PES complicated
- ▶ Bond-Length-Bond-Angle coordinates (Polyspherical Non-orthogonal coordinates)
  - ▶ Quartic PES and BC simple
  - ▶ KE complicated
- ▶ What is optimum?
  - ▶ BLBA: 15.43 cm<sup>-1</sup> error (1992)
  - ▶ Normal from Symmetric Jacobi: 10.79 cm<sup>-1</sup>
  - ▶ Radau: 1.49 cm<sup>-1</sup>
  - ▶ Optimized: 1.29 cm<sup>-1</sup>

---

<sup>1</sup>Kinetic Energy Operator

<sup>2</sup>Boundary Conditions

# Choosing coordinates: Step 1

- ▶ **X** matrix of nuclear Cartesians ( $\vec{X}$ )
  - ▶  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$

# Choosing coordinates: Step 1

- ▶ **X** matrix of nuclear Cartesians ( $\vec{X}$ )
  - ▶  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$
- ▶ **x** matrix of internal Cartesians ( $\vec{x}$ )
  - ▶ Normal coordinates:  $\vec{x} = \mathbf{L}\vec{X}$
  - ▶ Generalized coordinates:  $\mathbf{x} = \mathbf{X}\mathbf{M}$

# Choosing coordinates: Step 1

- ▶ **X** matrix of nuclear Cartesians ( $\vec{X}$ )
  - ▶  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$
- ▶ **x** matrix of internal Cartesians ( $\vec{x}$ )
  - ▶ Normal coordinates:  $\vec{x} = \mathbf{L}\vec{X}$
  - ▶ Generalized coordinates:  $\mathbf{x} = \mathbf{X}\mathbf{M}$ 
    - ▶  $T = -\frac{\hbar^2}{2} \sum_{\beta\beta'} \frac{1}{\mu_{\beta\beta'}} \sum_i \frac{\partial^2}{\partial x_{i\beta} \partial x_{i\beta'}}$
    - ▶  $\frac{1}{\mu_{\beta\beta'}} = \sum_{\alpha} M_{\alpha\beta} M_{\alpha\beta'} \frac{1}{m_{\alpha}}$

# Choosing coordinates: Step 1

- ▶ **X** matrix of nuclear Cartesians ( $\vec{X}$ )
  - ▶  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$
- ▶ **x** matrix of internal Cartesians ( $\vec{x}$ )
  - ▶ Normal coordinates:  $\vec{x} = \mathbf{L}\vec{X}$
  - ▶ Generalized coordinates:  $\mathbf{x} = \mathbf{X}\mathbf{M}$ 
    - ▶  $T = -\frac{\hbar^2}{2} \sum_{\beta\beta'} \frac{1}{\mu_{\beta\beta'}} \sum_i \frac{\partial^2}{\partial x_{i\beta} \partial x_{i\beta'}}$
    - ▶  $\frac{1}{\mu_{\beta\beta'}} = \sum_{\alpha} M_{\alpha\beta} M_{\alpha\beta'} \frac{1}{m_{\alpha}}$
- ▶ Orthogonal coordinates:  $\frac{1}{\mu_{\beta\beta'}} = \delta_{\beta\beta'} \frac{1}{\mu_{\beta\beta}}$

# Role of $\mathbf{M}$

- ▶  $\mathbf{M}$  only enters in computation of  $\frac{1}{\mu_{\beta\beta'}}$  and
- ▶ evaluation of potential:  $\mathbf{X} = \mathbf{xM}^{-1}$

# Role of $\mathbf{M}$

- ▶  $\mathbf{M}$  only enters in computation of  $\frac{1}{\mu_{\beta\beta'}}$  and
- ▶ evaluation of potential:  $\mathbf{X} = \mathbf{x}\mathbf{M}^{-1}$
- ▶ For BLBA,  $(\mathbf{M}^{-1})_{\beta\alpha} = \pm 1, 0$

# Role of $\mathbf{M}$

- ▶  $\mathbf{M}$  only enters in computation of  $\frac{1}{\mu_{\beta\beta'}}$  and
- ▶ evaluation of potential:  $\mathbf{X} = \mathbf{xM}^{-1}$
- ▶ For BLBA,  $(\mathbf{M}^{-1})_{\beta\alpha} = \pm 1, 0$
- ▶  $\det \mathbf{M} \neq 0$ ,  $\beta = N$  c.m. and  $\frac{1}{\mu_{N\beta}} = 0$ ,  $\beta \neq N$ .

## Choosing coordinates: Step 2

►  $\mathbf{x} = \mathbf{A}^T (\alpha^{LB} \beta^{LB} \gamma^{LB}) \mathbf{x}^{bf}$

►  $\mathbf{x}^{bf} = \begin{pmatrix} -r_1 \sin(1-a)\chi & r_2 \sin a\chi & r_3 \sin \theta_3 \cos \phi_3 & \dots \\ 0 & 0 & r_3 \sin \theta_3 \sin \phi_3 & \dots \\ r_1 \cos(1-a)\chi & r_2 \cos a\chi & r_3 \cos \theta_3 & \dots \end{pmatrix}$

## Choosing coordinates: Step 2

- ▶  $\mathbf{x} = \mathbf{A}^T (\alpha^{LB} \beta^{LB} \gamma^{LB}) \mathbf{x}^{bf}$

- ▶  $\mathbf{x}^{bf} = \begin{pmatrix} -r_1 \sin(1-a)\chi & r_2 \sin a\chi & r_3 \sin \theta_3 \cos \phi_3 & \dots \\ 0 & 0 & r_3 \sin \theta_3 \sin \phi_3 & \dots \\ r_1 \cos(1-a)\chi & r_2 \cos a\chi & r_3 \cos \theta_3 & \dots \end{pmatrix}$

- ▶  $T = ???$

## Choosing coordinates: Step 2

- ▶  $\mathbf{x} = \mathbf{A}^T (\alpha^{LB} \beta^{LB} \gamma^{LB}) \mathbf{x}^{bf}$

- ▶  $\mathbf{x}^{bf} = \begin{pmatrix} -r_1 \sin(1-a)\chi & r_2 \sin a\chi & r_3 \sin \theta_3 \cos \phi_3 & \dots \\ 0 & 0 & r_3 \sin \theta_3 \sin \phi_3 & \dots \\ r_1 \cos(1-a)\chi & r_2 \cos a\chi & r_3 \cos \theta_3 & \dots \end{pmatrix}$

- ▶  $T = ???$

- ▶ Bramley and Handy tetra-atomic:  $T$  has 75 terms ...

## Choosing coordinates: Step 2

- ▶  $\mathbf{x} = \mathbf{A}^T (\alpha^{LB} \beta^{LB} \gamma^{LB}) \mathbf{x}^{bf}$

- ▶  $\mathbf{x}^{bf} = \begin{pmatrix} -r_1 \sin(1-a)\chi & r_2 \sin a\chi & r_3 \sin \theta_3 \cos \phi_3 & \dots \\ 0 & 0 & r_3 \sin \theta_3 \sin \phi_3 & \dots \\ r_1 \cos(1-a)\chi & r_2 \cos a\chi & r_3 \cos \theta_3 & \dots \end{pmatrix}$

- ▶  $T = ???$

- ▶ Bramley and Handy tetra-atomic:  $T$  has 75 terms ...
- ▶ basis functions: anything with finite matrix elements

# Corrections to Born-Oppenheimer Approximation (2001)

- ▶ B.O.:  $V$  from  $H^{el}\psi^{el} = E\psi^{el}$ ,  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$

# Corrections to Born-Oppenheimer Approximation (2001)

- ▶ B.O.:  $V$  from  $H^{el}\psi^{el} = E\psi^{el}$ ,  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$
- ▶ B.O.:  $V$  from  $H^{el}\psi^{el} = E\psi^{el}$ ,  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$
- ▶ 1<sup>st</sup> order:  $V$  from  $\langle \psi^{el} | H^{el} + T | \psi^{el} \rangle$ ,  $T = \text{same}$

# Corrections to Born-Oppenheimer Approximation (2001)

- ▶ B.O.:  $V$  from  $H^{el}\psi^{el} = E\psi^{el}$ ,  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$
- ▶ B.O.:  $V$  from  $H^{el}\psi^{el} = E\psi^{el}$ ,  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$
- ▶ 1<sup>st</sup> order:  $V$  from  $\langle \psi^{el} | H^{el} + T | \psi^{el} \rangle$ ,  $T = \text{same}$
- ▶ 2<sup>nd</sup> order:  $\psi^{el}$  changes, so  $V = \text{same} + C^{(0)}$ ,  
 $T = \text{same} + \sum_{i\alpha i'\alpha'} C_{i\alpha i'\alpha'}^{(2)} \frac{\partial^2}{\partial X_{i\alpha} \partial X_{i'\alpha'}} + \sum_{i\alpha} C_{i\alpha}^{(1)} \frac{\partial}{\partial X_{i\alpha}}$

# Easier way to derive $T$

$$\blacktriangleright T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$$

## Easier way to derive $T$

- ▶  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$
- ▶  $= +\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \left( \frac{\partial}{\partial X_{i\alpha}} \right)^{\dagger} \frac{\partial}{\partial X_{i\alpha}}$

# Easier way to derive $T$

- ▶  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$
- ▶  $= +\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \left( \frac{\partial}{\partial X_{i\alpha}} \right)^{\dagger} \frac{\partial}{\partial X_{i\alpha}}$ 
  - ▶ bi-linear form

# Easier way to derive $T$

- ▶  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$
- ▶  $= +\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \left( \frac{\partial}{\partial X_{i\alpha}} \right)^{\dagger} \frac{\partial}{\partial X_{i\alpha}}$ 
  - ▶ bi-linear form
  - ▶  $\nabla^2 = \frac{1}{r} \frac{\partial^2}{\partial r^2} r + \frac{1}{r^2} \left( \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \sin \theta \frac{\partial}{\partial \theta} + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2} \right)$

# Easier way to derive $T$

- ▶  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$
- ▶  $= +\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \left( \frac{\partial}{\partial X_{i\alpha}} \right)^{\dagger} \frac{\partial}{\partial X_{i\alpha}}$ 
  - ▶ bi-linear form
  - ▶  $\nabla^2 = \frac{1}{r} \frac{\partial^2}{\partial r^2} r + \frac{1}{r^2} \left( \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \sin \theta \frac{\partial}{\partial \theta} + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2} \right)$
  - ▶  $= -\left( \frac{\partial}{\partial r} \right)^{\dagger} \frac{\partial}{\partial r} - \left( \frac{\partial}{\partial \theta} \right)^{\dagger} \frac{1}{r^2} \frac{\partial}{\partial \theta} - \left( \frac{\partial}{\partial \phi} \right)^{\dagger} \frac{1}{r^2 \sin^2 \theta} \frac{\partial}{\partial \phi}$

# Easier way to derive $T$

- ▶  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$
- ▶  $= +\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \left( \frac{\partial}{\partial X_{i\alpha}} \right)^{\dagger} \frac{\partial}{\partial X_{i\alpha}}$ 
  - ▶ bi-linear form
- ▶  $= +\frac{\hbar^2}{2} \sum_{\beta\beta'} \frac{1}{\mu_{\beta\beta'}} \sum_i \left( \frac{\partial}{\partial x_{i\beta}} \right)^{\dagger} \frac{\partial}{\partial x_{i\beta'}}$

# Easier way to derive $T$

- ▶  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$
- ▶  $= +\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \left( \frac{\partial}{\partial X_{i\alpha}} \right)^{\dagger} \frac{\partial}{\partial X_{i\alpha}}$ 
  - ▶ bi-linear form
- ▶  $= +\frac{\hbar^2}{2} \sum_{\beta\beta'} \frac{1}{\mu_{\beta\beta'}} \sum_i \left( \frac{\partial}{\partial x_{i\beta}} \right)^{\dagger} \frac{\partial}{\partial x_{i\beta'}}$
- ▶  $\frac{\partial}{\partial x_{i\beta}} = \sum_j A_{ji} (\alpha^{LB} \beta^{LB} \gamma^{LB}) \sum_l g_{j\beta l} \hat{O}_l$ 
  - ▶  $\hat{O}_l = \frac{\partial}{\partial r_1}, \frac{\partial}{\partial r_2}, \frac{\partial}{\partial \chi}, \dots, \hat{J}_x, \hat{J}_y, \text{ and } \hat{J}_z.$


# Easier way to derive $T$

- ▶  $T = -\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \frac{\partial^2}{\partial X_{i\alpha}^2}$
- ▶  $= +\frac{\hbar^2}{2} \sum_{\alpha} \frac{1}{m_{\alpha}} \sum_i \left( \frac{\partial}{\partial X_{i\alpha}} \right)^{\dagger} \frac{\partial}{\partial X_{i\alpha}}$ 
  - ▶ bi-linear form
- ▶  $= +\frac{\hbar^2}{2} \sum_{\beta\beta'} \frac{1}{\mu_{\beta\beta'}} \sum_i \left( \frac{\partial}{\partial x_{i\beta}} \right)^{\dagger} \frac{\partial}{\partial x_{i\beta'}}$
- ▶  $\frac{\partial}{\partial x_{i\beta}} = \sum_j A_{ji} (\alpha^{LB} \beta^{LB} \gamma^{LB}) \sum_l g_{j\beta l} \hat{O}_l$ 
  - ▶  $\hat{O}_l = \frac{\partial}{\partial r_1}, \frac{\partial}{\partial r_2}, \frac{\partial}{\partial \chi}, \dots, \hat{J}_x, \hat{J}_y, \text{ and } \hat{J}_z.$
- ▶ for openshell systems ...

# What If All Orthogonal Coordinates Not Very Good?


# What If All Orthogonal Coordinates Not Very Good?

► SO<sub>2</sub>



# What If All Orthogonal Coordinates Not Very Good?

- $\text{SO}_2$
- $\text{C}_3$



# What If All Orthogonal Coordinates Not Very Good?

- ▶  $\text{SO}_2$
- ▶  $\text{C}_3$
- ▶  $\text{C}_3\text{H}^-$

# How to optimize **M**

- Constraints:  $\beta = N$  c.m.,  $\frac{1}{\mu_{\beta N}} = 0$

# How to optimize **M**

- ▶ Constraints:  $\beta = N$  c.m.,  $\frac{1}{\mu_{\beta N}} = 0$
- ▶  $\mu_{\beta\beta} = 1$

# How to optimize **M**

- ▶ Constraints:  $\beta = N$  c.m.,  $\frac{1}{\mu_{\beta N}} = 0$
- ▶  $\mu_{\beta\beta} = 1$
- ▶  $V$  cross terms small and  $\frac{1}{\mu_{\beta\beta'}}$  small,  $\beta \neq \beta'$

# How to optimize **M**

- ▶ Constraints:  $\beta = N$  c.m.,  $\frac{1}{\mu_{\beta N}} = 0$
- ▶  $\mu_{\beta\beta} = 1$
- ▶  $V$  cross terms small and  $\frac{1}{\mu_{\beta\beta'}}$  small,  $\beta \neq \beta'$
- ▶ All  $r_e$  roughly the same

# How to optimize **M**

- ▶ Constraints:  $\beta = N$  c.m.,  $\frac{1}{\mu_{\beta N}} = 0$
- ▶  $\mu_{\beta\beta} = 1$
- ▶  $V$  cross terms small and  $\frac{1}{\mu_{\beta\beta'}}$  small,  $\beta \neq \beta'$
- ▶ All  $r_e$  roughly the same
- ▶ Repeatedly guess and walk downhill

# Summary

- ▶ 1980's Handy introduces Bond-length-bond-angle coordinates

# Summary

- ▶ 1980's Handy introduces Bond-length-bond-angle coordinates
- ▶ 1992 I scoff at BLBA coordinates

# Summary

- ▶ 1980's Handy introduces Bond-length-bond-angle coordinates
- ▶ 1992 I scoff at BLBA coordinates
- ▶ 2001 I use BLBA coordinates

# Summary

- ▶ 1980's Handy introduces Bond-length-bond-angle coordinates
- ▶ 1992 I scoff at BLBA coordinates
- ▶ 2001 I use BLBA coordinates
- ▶ today I embrace generalized coordinates